

ROLE OF THE HEALTH DEPARTMENT

State and local health agencies monitor plague activity throughout the State. Rangers, park personnel and others are trained to watch for sick or dead rodents or other evidence that plague may be active in a particular area and to report their findings to health authorities.

Health authorities will institute preventative measures when animal plague is found in areas with human exposure. Warning signs will be posted. After careful evaluation, the area may be quarantined and insecticides may be used to reduce the risk of flea bites to humans.

Insecticide dust is applied into rodent burrows and/or into tube-like containers called "bait stations." Rodents enter the bait stations and get flea powder in their fur. They also carry the insecticide in their fur back to the nest, killing fleas inside the burrows. This method of flea control is very effective, uses a minimum of insecticide, and does not harm the rodents.

IF YOU SEE A BAIT STATION, PLEASE DO NOT DISTURB IT.

Additional information on plague and other vector-borne diseases can be obtained from your local vector control district or local health department. For more information call:

MOSQUITO and VECTOR MANAGEMENT DISTRICT of Santa Barbara County

OFFICE: 2450 Lillie Ave. * MAIL: P.O. Box 1389
Summerland, CA 93067
TELEPHONE: (805) 969-5050
E-MAIL: mvmdistrict@mvmdistrict.org
Visit our WEBSITE: www.mvmdistrict.org

This brochure was developed by the State of California, Department of Public Health, Vector-Borne Disease Section. (916) 552-9730.

FACTS ABOUT PLAGUE IN CALIFORNIA




Plague is a highly infectious bacterial disease which primarily affects rodents. Humans and their pets (dogs, and especially cats) can get plague if they visit or live in areas where wild rodents are naturally infected. The disease may also occur in rats that can live in close contact with people.

This is the same disease that ravaged Europe in the 6th Century (the Plague of Justinian) and again in the 14th Century (the Black Death). At the present time, plague in humans is relatively rare, and can be treated successfully with modern antibiotics. However, it is vital that the disease be diagnosed and treated in its early stages. If not, it is often fatal and, if lung infection (plague pneumonia) develops, it can be transmitted directly and rapidly to others.

EXPOSURE TO PLAGUE

People can get the disease from animals in several ways. The most important routes of transmission are:

1. BITES OF FLEAS FROM INFECTED RODENTS.




Hungry fleas will leave a sick or dead rodent to find another host and can bite people.

2. DIRECT CONTACT WITH SICK ANIMALS.




The bacteria in the blood or other body fluids of an infected animal can enter through cuts and abrasions on the hands or mucous membranes.

3. PET INVOLVEMENT. (1) Infected rodent fleas can be brought into the home or campsite by a dog or cat. (2) Plague pneumonia can be caught from a sick cat that is coughing or sneezing.



RISKS OF PLAGUE

Plague in California occurs in the foothills, plateaus, mountains, and foggy coastal belt (shaded areas on map). Plague is absent from the southeastern desert region and the Central Valley.



The major threat of plague to humans is in the recreational, wilderness, and rural parts of the State as well as the suburban foothill areas of some of our large metropolitan cities.


Plague is most often associated with wild rodents, however, it may be transferred to rats in urban areas, increasing the risk of plague transmission to humans. The last urban outbreak occurred in Los Angeles in 1924 and 1925.

Due to the potential danger of pneumonic plague epidemics, public health authorities place a high priority on prevention of human plague cases. You can minimize your exposure to plague by carefully following the precautions listed here.

WHICH ANIMALS CARRY PLAGUE?



Chipmunk



Golden-Mantled Ground Squirrel

The most important wild rodents that can carry the disease are squirrels (especially ground squirrels), chipmunks, wood rats, mice, and marmots. Plague is lethal to many rodents, therefore, any sign of sick or dead rodents is a warning that plague may be in the area. Other wild animals -- especially rabbits, carnivores (including coyote, bobcat, badger, bear, gray fox, and skunk), and wild pigs -- can also acquire plague, but usually with no signs of illness.

Domestic animals can acquire plague and pose a direct threat to humans. Dogs rarely become ill, but cats are highly susceptible and can suffer a severe illness. Pets can transport rodent fleas from the field into homes or campsites. If a cat develops plague pneumonia, it can infect humans by coughing and sneezing.

Ways to Protect Your Family From Plague

General Precautions

- ▶ If you become ill within seven days after possible exposure to plague, contact a physician immediately.
- ▶ Use caution when handling a sick pet that has been in a plague area, especially a cat. Avoid face-to-face contact. Consult a veterinarian. Inform the vet that the animal has been in a plague area.
- ▶ Avoid all contact with rodents and their fleas. Do not touch sick or dead rodents. Report them to rangers or health authorities.

Where You Live

- ▶ Keep rodent populations down around homes and other inhabited areas. Prevent them from entering buildings. As much as possible, remove or deny rodents access to any source of food or shelter.
- ▶ Minimize pet contact with rodents and their fleas. A bell on a cat's collar may help. Protect pets with flea powder. Flea collars are helpful, but, if used alone, are too slow acting.

Where You Work or Play

- ▶ Do not camp, sleep, or rest near animal burrows.
- ▶ Do not feed rodents or other wild animals in campgrounds and picnic areas. Store food and refuse in rodent-proof containers.
- ▶ Wearing long pants tucked into boot tops can reduce your exposure to fleas. Insect repellent sprayed on socks and trouser cuffs also may help.
- ▶ LEAVE PETS AT HOME if possible. If not, keep pet confined or on a leash. Do not allow pet to approach sick or dead rodents or to explore rodent burrows. Protect pet with flea powder (flea collars alone are too slow acting).
- ▶ HUNTERS AND TRAPPERS: Use rubber gloves when skinning and cleaning rodents, rabbits, wild pigs, coyotes and other carnivores. Cook these animals thoroughly.

SYMPTOMS OF PLAGUE

In humans, the initial symptoms of plague include fever, chills, muscle aches, a feeling of weakness and, commonly, swollen and tender lymph nodes (called "buboes"). This stage is called bubonic plague.

Contact a Physician Immediately If You Become Ill Within 7 Days of Being in a Plague Area.

The usual incubation period is 2 to 6 days. Plague is curable when diagnosed early. You can help with the diagnosis by telling your doctor where you have been and what you have done that may have exposed you to plague.

If it is not treated in time, bubonic plague can progress to septicemic plague (bloodstream infection) and/or pneumonic plague (plague pneumonia).

A cat with plague will become very ill, may stop eating, and will have a fever. Swollen lymph nodes may occur, generally in the neck area.

Inform the Veterinarian If a Sick Pet Has Been in a Plague Area.

